

**GAVISCON INFANT**  
**Magnesium alginate, Sodium alginate**  
**Powder for Oral Suspension**  
**PATIENT INFORMATION LEAFLET**

**Please read all of this leaflet carefully because it contains important information for you and your child.**

This medicine is available without prescription. However, you still need to use Gaviscon Infant carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- You must contact a doctor if your child's symptoms worsen or do not improve after 7 days.
- If any of the side effects get serious, or if you notice any side effect not listed in this leaflet, please tell your doctor or pharmacist.

**In this leaflet:**

1. What Gaviscon Infant is and what it is used for
2. Before using Gaviscon Infant
3. How to use Gaviscon Infant
4. Possible side effects
5. How to store Gaviscon Infant
6. Further Information

**FPO**

**1. What Gaviscon Infant is and what is it used for?**

Gaviscon Infant contains sodium alginate and magnesium alginate which belong to a group of medicines called "reflux suppressants".

Reflux is the process by which the stomach contents flow back up the tube that leads from the mouth to the stomach (known as the gullet or oesophagus). This can cause an infant to be sick, spit up more than normal and become distressed. Gaviscon Infant prevents reflux by stabilising the stomach contents, so reducing the incidence of reflux and bringing up of feeds (possetting).

**2. Before using Gaviscon Infant**

**Do not use Gaviscon Infant:**

- if your baby was born pre-maturely
- if your child is allergic to magnesium alginate, sodium alginate or any of the other ingredients (see section 6)
- if your child is being sick, has a fever or diarrhoea
- if you have been told or it is suspected that your child has damage to their kidneys or they have been put on a low salt diet
- if you have been told that your child has a blockage in their gut (an intestinal obstruction)

**Take special care with Gaviscon Infant:**

If your baby is under one year, you should ask your doctor or pharmacist before giving Gaviscon Infant.

**Taking Gaviscon Infant with other medicines**

Please tell your doctor or pharmacist if your child is taking or has recently taken any other medicines, including medicines obtained without a prescription.

**Taking Gaviscon Infant with food and drink**

**Do not** use Gaviscon Infant with other feed thickening agents or infant milk preparations containing a thickening agent. You should ask your doctor or pharmacist before you give Gaviscon Infant to your child if you are not sure.

**Important information about some of the ingredients of Gaviscon Infant**

This medicinal product contains 0,9 mmol (or 21 mg) sodium per sachet. To be taken into account by patients on a controlled sodium diet.

Gaviscon Infant should not be given to infants with kidney problems/damage or those on a low salt diet (see 'Do not use Gaviscon Infant').

### **3. How to use Gaviscon Infant**

For infants aged 1 to 2 years. Not to be used in premature infants or infants under one year except under medical supervision. Not suitable for children over 2 years, adults or the elderly.

Always use Gaviscon Infant exactly as described below. You should check with your doctor or pharmacist if you are not sure.

The dosage and mixing instructions depend on the weight of your child and the feeding method you use:

Infants under 4.5kg (10lb) – one sachet should be used

Infants over 4.5kg (10lb) – two sachets should be used

Open the sachet(s) and mix the contents immediately before use as directed below:

Bottle fed infants:	<ul style="list-style-type: none"><li>• Mix each sachet into 115ml (4 fluid oz) of feed in the bottle</li><li>• Shake well</li><li>• Feed as normal</li></ul>
Breast fed infants and other infants up to 2 years of age:	<ul style="list-style-type: none"><li>• Mix each sachet with 5ml (1 teaspoon) of cooled, boiled water until a smooth paste is formed</li><li>• Add another 10ml (2 teaspoons) of cooled, boiled water and mix</li><li>• For breast fed infants give Gaviscon Infant part way through each feed or meal using a spoon or feeding bottle</li><li>• For all other infants give Gaviscon Infant at the end of each meal using a spoon or feeding bottle.</li></ul>

Do not give more than six times in 24 hours.

**If your child's condition persists or becomes worse ask your doctor or other professional for advice immediately.**

You **should not** use Gaviscon Infant for more than seven days without checking with your doctor or other health professional.

**If your child takes more Gaviscon Infant than they should:**

If your child takes too much Gaviscon Infant they may be sick, lose their appetite, become irritable or tired and weak.

If you accidentally give too much Gaviscon Infant ask your doctor or other health professional for advice **immediately**.

**If you forget to give Gaviscon Infant:**

If a dose is missed **do not** give a double dose with the next feed. Just carry on with the recommended dose.

If you have any further questions on the use of this product ask your doctor or pharmacist.

### **4. Possible side effects**

Like all medicines, Gaviscon Infant can cause side effects, although not everybody gets them.

Very rarely, Gaviscon Infant can cause bloating and discomfort of the stomach (gastric distension).

If the child has any side effects, stop giving Gaviscon Infant and tell your doctor or other healthcare professional.

### **5. How to store Gaviscon Infant**

**Keep all medicines out of reach and sight of children.**

Do not use after the expiry date shown on the pack (EXP month/year).

Do not store above 30°C.

Do not use if the sachet is damaged or the security seal is broken.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

### **6. Further information**

Each single dose sachet of Gaviscon Infant contains 225 mg sodium alginate and 87.5 mg magnesium alginate as the active ingredients.


The other ingredients are mannitol (E421) and colloidal silica.

**What Gaviscon Infant looks like and the contents of the pack**

Gaviscon Infant is a powder. Each carton contains 30 single dose sachets of powder, joined in pairs.

**Marketing Authorisation Holder and Manufacturer**

Reckitt Benckiser Healthcare (UK) Limited, Dansom Lane, Hull, HU8 7DS.

Gaviscon and  are trademarks      Leaflet last revised: September 2012