

Robitussin[®]

CHESTY COUGH MEDICINE

PAA129413

Guaifenesin 100 mg per 5 ml Liquid for Oral Administration

Read all of this leaflet carefully because it contains important information for you

This medicine is available without prescription. However, you still need to use this medicine carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- You must contact a doctor if your symptoms worsen or do not improve after 7 days.
- If any of the side effects get serious, or if you notice any side effect not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Robitussin Chesty Cough Medicine is and what it is used for
2. Before you take Robitussin Chesty Cough Medicine
3. How to take Robitussin Chesty Cough Medicine
4. Possible side effects
5. How to store Robitussin Chesty Cough Medicine
6. Further information

1. What Robitussin Chesty Cough Medicine is and what it is used for

This medicine contains the active ingredient guaifenesin. This belongs to a group of medicines called cough expectorants that work by changing an unproductive cough to a cough that is more productive and less frequent.

2. Before you take Robitussin Chesty Cough Medicine

Do not take this medicine if:

- You are **allergic** (hypersensitive) to **guaifenesin** or any of the other ingredients of this medicine (see section 6 for ingredient list).

Talk to your doctor before taking this medicine if:

- You have had a cough for a long time or if you have **asthma** or are suffering from an **asthma attack**.

Pregnancy and breastfeeding:

Please talk to your doctor before taking this medicine if you are pregnant or breastfeeding.

Important information about some of the ingredients in this medicine:

- If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine as it contains **sorbitol** and **maltitol**.
- This medicinal product contains **2.7 % v/v alcohol** (ethanol) (214 mg per 10 ml dose). This is equivalent to approximately 2 ml wine per dose which can be harmful for those suffering from alcoholism and should be taken into account if you are pregnant or breastfeeding or have liver disease or epilepsy. This will have no or negligible influence on your ability to drive.

3. How to take Robitussin Chesty Cough Medicine

For oral use only.

Adults, the elderly and children over 12 years: Take 10 ml up to four times daily, as required.

Children under 12 years: Do not use.

Use the dosing cup provided with the pack.

Do not exceed the stated dose. No more than four doses should be taken in any 24 hours.

Stop using this medicine and ask a healthcare professional if you have had a cough for **more than 7 days** or if you have any other symptoms which might be related to your cough, for example fever, rash, persistent headache or shortness of breath.

If you take more medicine than you should:

If you take too much of this medicine, consult your doctor or hospital immediately. Bring any remaining medicine with you to show the doctor.

If you forget to take your medicine:

Take a dose when you need to, provided the last dose was taken at least 4 hours ago. Do not take a double dose to make up for the one you have missed.



4. Possible side effects

This medicine can cause side effects, although not everybody gets them. Side effects with guaifenesin are rare. However, some susceptible individuals may experience unwanted side effects.

If you experience any of the following side effects mentioned below or any other side effects, **STOP** taking this medicine immediately and contact your doctor:

- Allergic reactions. Symptoms could include itchy skin or rashes, difficulty breathing, faster heart rate, swelling of the face or throat.

Other possible side effects:

- Feeling sick (nausea) or being sick (vomiting).

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Robitussin Chesty Cough Medicine

- Do not use this medicine after the expiry date shown on the top end of the carton.
- Do not store above 25°C.
- Keep out of the sight and reach of children.

6. Further information

What Robitussin Chesty Cough Medicine contains:

Each 5 ml contains 100 mg of the active substance guaifenesin.

Robitussin Chesty Cough also contains: Glycerol, Carmellose Sodium, Sodium Benzoate (E211), Sodium Cyclamate, Ethanol, Levomenthol, Maltitol (E965), Sorbitol Solution (70%), Natural Cherry Flavour, Citric Acid Anhydrous, Caramel (E150), Acesulfame Potassium, Purified Water.

Robitussin Chesty Cough Medicine is a liquid, supplied in a 100 ml bottle or 250 ml bottle. Not all pack sizes maybe marketed.

Who makes and supplies this medicine:

The Marketing Authorisation holder for Robitussin Chesty Cough Medicine is: Pfizer Consumer Healthcare Ltd, Ramsgate Road, Sandwich, Kent, CT13 9NJ, UK.

The dosing cup is manufactured by:

Humphrey Cosburn Plastics, 89 Union Street, Toronto, Ontario, M6N 3N2, Canada.

The EU Authorised Representative is: Pfizer Healthcare Ireland

This medicine is manufactured by:

Wyeth Pharmaceuticals, New Lane, Havant, Hampshire, PO9 2NG, United Kingdom.
Pfizer Consumer Manufacturing Italy S.r.l., Via Nettunense, 90, 04011 Aprilia (LT), Italy.

If you have any queries, comments about this medicine or require information in large print, tape or CD please contact the Careline by phone or write to:

Careline, Pfizer Consumer Healthcare Ltd,
Walton Oaks, Surrey, KT20 7NS, UK.
or carelineUK@pfizer.com



Leaflet prepared: May 2019
PL 00165/0097
MA111/00403

PAA129413

