


CARRIER



LEAFLET FACE

92.0 mm	<p>5- How to store this medicine KEEP OUT OF THE REACH AND SIGHT OF CHILDREN. Store below 25°C. Use within 14 days after first opening. Use before the expiry date shown on the front of the bottle.</p> <p>6- Further information Woodward's Gripe Water - Alcohol Free & Sugar Free is an oral solution containing Terpeneless Dill Seed Oil 2.3mg/5mL, and Sodium Hydrogen Carbonate 52.5mg/5mL as active ingredients and also Maltitol Liquid, Glycerol, Sodium Salts of Esters of p-Hydroxybenzoic Acid (E215, E217 and E219), Disodium Edetate, Propylene Glycol and Purified Water.</p>	<p>Licence Holder: Reckitt Benckiser Healthcare (UK) Ltd, 103-105 Bath Road Slough SL1 3UH Tel 0500 455 456</p> <p>Manufacturer : SSL International Plc, Peterlee, SR8 2RU. FAMAR A.V.E, 49th National Road Athens-Lamia, Avlona Attiki, 19011 Greece</p> <p>PL 00063/0698</p> <p>Last leaflet update 06/2013</p>	<p>Read the leaflet carefully before use and for full list of ingredients.</p> <p>Store below 25°C. Use within 14 days of first opening the bottle.</p>	
	Page 8,9	Page 10,11	Page 12	Page 1
	60.00 mm	60.00 mm	30.00 mm	33.00 mm

LEAFLET REVERSE

92.0 mm	<p>Patient Information Woodward's Gripe Water - Alcohol Free & Sugar Free Terpeneless Dill Seed Oil 2.3mg/5mL, Sodium Hydrogen Carbonate 52.5mg/5mL</p> <p>1- What is this medicine used for This medicine is an antacid and antifatulent product. It is used to relieve griping pain or waves of severe abdominal pain (infant colic) in babies, due to trapped air in the tummy. It can be given to babies from 1 to 12 months.</p> <p>2- Before you use this medicine Do not give to a baby: - who is allergic to any of the ingredients in this product (see section 6) - who has kidney problems.</p>	<p>Important information about some of the ingredients This product is sugar free but contains maltitol, which may affect blood sugar levels. Therefore if your baby has an inherited intolerance to some sugars, check with your doctor before giving this medicine.</p> <p>This medicine contains hydroxybenzoate derivatives (E215, E217 and E219) which may cause allergic reactions (possibly delayed).</p> <p>This medicinal product contains 0.7 mmol (or 16.1 mg) sodium per 5mL dose and 1.4 mmol (32.2 mg) per 10mL dose.</p>	<p>3- How to use this medicine This medicine is only for babies from 1 to 12 months old. 1-6 months old: one 5mL spoonful 6-12 months old: two 5mL spoonfuls Give during or after a feed. Do not use more than 6 times in 24 hours. If symptoms persist, contact your doctor. If you give your baby too much of this medicine, speak to a doctor or pharmacist at once.</p> <p>4- Possible side effects If your baby experiences any side effect after using this product, stop using it and seek medical advice.</p>
	Page 2,3	Page 4,5	Page 6,7
	63.00 mm	60.00 mm	60.00 mm